

# ESA update for CEOS LSI-VC

**Susanne Mecklenburg**  
European Space Agency  
Sentinel-3 and SMOS Mission Manager

LSI-VC-5, Tokyo, 21-23 February 2018

**Thanks to ESA colleagues  
for contributions:**


F.Gascon, P.Potin, B.Rosich,  
F.Niro, S.Dransfeld,  
N.Miranda, N.Mileva, and  
S.Hosford (CNES)

- Update on Copernicus Space Segment
- ESA's contribution to FDA activities
  - Copernicus Data Access
  - Data and Information Access Services (DIAS)
- ESA's contribution to ARD activities
  - Current data products over land
  - Future developments
- Further ARD consultations
  - Land Product Validation and Evolution Workshop
  - IGARSS 2018

# Copernicus Space Segment – update

# Sentinel-1 overall mission status

- Sentinel-1 **nominal routine operations on-going**
  - Overall mission in a very good shape
  - The mission provides: global and routine coverage, with a systematic production scenario,
  - Sentinel-1A and -1B data routinely provided to Copernicus Services and users worldwide
  - On-going support to various activations from the Copernicus Emergency Management Service and International Charter Space and Major Disasters
- **Sentinel-1 constellation generates now 11 TB of products daily** (against a specification of 3 TB)
- **Sentinel-1 is operated close to its full mission capacity** (i.e. difficulty to accommodate additional observations)
- **Upcoming activities**
  - Identify and implement mission evolution to support new user needs, assess an increased system capacity enabling further mission exploitation, etc.



Md 4.0 Earthquake in Ischia, Italy, 21/08/2017.

Ground deformation measured by Sentinel-1


© Contains Copernicus Sentinel Data (2017) / CNR-IREA



# Sentinel-1 Constellation Observation Scenario: Mode - Polarisation - Observation Geometry



validity start: 05/2017



## POLARISATION SCHEMA

HH or HH-HV

VV or VV-HV

HH or HH-HV

## MODE / POLARISATION

- IW mode / dual polarisation
- IW mode / single polarisation
- EW mode / dual polarisation
- EW mode / single polarisation

- SM mode / dual-polarisation
- SM mode / single-polarisation

- Calibration Site  
(locally different modes or polarisations possible)


## PASS

- ASCENDING
- DESCENDING

# Sentinel-1 Constellation Observation Scenario: Revisit & Coverage Frequency



validity start: 05/2017



## PASS

- ASCENDING
- DESCENDING

## REVISIT

- 6 days
- 12 days

## FREQUENCY \*

- 12 days
- 24 days

## COVERAGE

- 1 days
- 1-3 days
- 2-4 days

## FREQUENCY \*\*

- 1 days
- 1-3 days
- 2-4 days

## REFERENCE DATA SITES (6d repeat)


- Highly active volcanism
- Fast subsidence
- Short growth cycle, intensive agriculture
- Fast changing wetlands
- Fast moving outlet glaciers
- Permafrost & glaciers

\* coverage ensured from same, repetitive relative orbits  
 \*\* coverage not considering repetitiveness of relative orbits


# Sentinel-2 mission status

- Sentinel-2A and -2B are together acquiring Europe, Africa and Greenland with 5-day revisit and the rest of the World is revisited every 6.6 days in average. Global systematic 5-day revisit will be reached once EDRS or the 4th ground station (Inuvik) become fully operational.
- In addition to the baseline acquisition areas, Antarctica is being imaged every 10 days.
- A new version of the Sentinel-2A spectral response functions has been released. These updated Sentinel-2A spectral responses are mainly changing the responses for bands B01 and B02, correcting inaccurate spectral responses provided previously.
- The systematic global production of Level 2A core products is being implemented in the ground segment. Systematic production using Sen2Cor algorithm is planned to start in Q1 2018 for Europe, and reach full global coverage in mid 2018.

**Sentinel-2 Constellation Observation Scenario:  
Revisit Frequency**










# Sentinel-3 mission status

- Sentinel-3A is now in **routine operations phase** and has reached full operational capacity: ESA-EUMETSAT Joint Routine Operations Readiness Review – successfully completed in October 2017.
- Overall status of Sentinel-3A is **nominal**, with all space and ground subsystems and instruments (OLCI, SLSTR, SRAL, MWR) performing nominally.
- **All Level 1 and Level 2 Land and Marine core data product have been released.**
- Definition and implementation of **new core data products** (AOD and FRP, EC change request) on-going; sample products expected in Q1/Q2 2018
- **Reprocessing** campaigns, including data from commissioning phase are ongoing for optical and altimetry data.
- Intense on ground **validation activities** are on-going to ensure the best possible user acceptance of the Sentinel-3A products.
- Preparations for **Sentinel-3B launch, planned for April 2018, on-going**, including implementation of the Tandem Phase and orbit phase shift to 140 degree between Sentinel-3A and –B



**Land Surface Temperature monthly composite for September 2016; Credit: D. Ghent, University of Leicester**

## Sentinel-1

- ★ With the authorization of the Satellite Production Review (SPR) Board, Sentinel-1C and -1D procurement programme progressed well with the production of flight units and with the successful closure of the AIS Receiver PDR (Preliminary Design Review).
- ★ The satellite platform, the SAR instrument and the Optical Communication Payload (OCP) are progressing according to plan.

## Sentinel-2

- ★ The production of flight units progressed well at platform and instrument level. With regards to the Sentinel-2C/-2D platform procurements, all delivery dates are still in line with the project need dates.
- ★ The Multi-Spectral Instrument (MSI) equipment manufacturing is progressing as planned, except for the SOFRADIR SWIR detectors. Mitigation actions are in place to reduce schedule impact.

# CSC Development Status: Space Segment

## Sentinel-3

- ★ The Sentinel-3B satellite is fully integrated with all flight units. The Assembly, Integration and Test Phase has been completed in Quarter 4/2017, which comprised platform tests, System Validation Test and launch adapter fit checks for both Rockot and Vega.
- ★ The Sentinel-3B Flight Acceptance Review (FAR) closed successfully with the Board meeting on 12 December 2017.
- ★ Following the request from EC to maintain the launch activities with Eurockot as per nominal launch service contract, the Agency proceeded with the launch preparations for Rockot. The present Sentinel-3B launch slot is from 30 March to 30 April 2018, with tentative launch date starting from mid April 2018 (tbc).
- ★ The Sentinel-3C and -3D development activities are proceeding as planned at platform/instruments level. The Sea and Land Surface Temperature Radiometer (SLSTR) and the Ocean and Land Colour Instrument (OLCI) remain the main drivers of the Sentinel-3C schedule, as such followed-up closely by the project team.



**Sentinel-3B during the Rockot  
Launcher Fit Check Test**

# A Long Term Scenario (LTS)



- Fundamental aspects of a LTS:
  - **assure user-driven continuity** and increase the **robustness** of the existing CSC in the future (Priority)
  - **increase the quality and quantity** of the existing measurements
  - **expand observation** types according to policies and user needs
  - employ **latest technologies** for maximum efficiency
  - **Partnerships and cooperation** are essential to success
- **Key driver is the evolving needs of the services prioritized by EC** through various consultative processes over the last year



ESA UNCLASSIFIED - For Official Use

Susanne de 12



European Space Agency



- ★ **November 2017 => Draft CSC Long-Term Scenario** submitted to ESA's Earth Observation Programme Board
- ★ **December 2017 => ESA released the Invitation To Tenders for Copernicus Space Component Phases A/B1 of Potential Expansion Sentinels:**
  - ★ Phase A/B1 of Anthropogenic CO2 Monitoring Mission
  - ★ Phase A/B1 of High Spatio-Temporal Resolution Land Surface Temperature (LST) Monitoring Mission
  - ★ Phase A/B1 of Polar Ice and Snow Topographic Mission
  - ★ Phase A/B1 of Passive Microwave Imaging Mission
  - ★ Phase A/B1 of HyperSpectral Imaging Mission
  - ★ Phase A/B1 L-Band SAR Mission
- ★ Activities related to the above are expected to start by mid 2018.


## ESA FDA activities – Data Access

## 4 Sentinel data access hubs operated by ESA




**Statistics at end December 2017**

# Sentinels Data Access at ESA - Statistics




*Some evidences of the success of Copernicus:*

1. A continuous increase of users registered on Open Access Hub (+ 100% in 1 year)
2. A weekly average of 0.7 PB of data downloaded by users



**Total volume of data downloads during last 3 months → 8.3 PBytes**






# Sentinels Data Access – Open Access Hub



Distribution of active users on ESA  
Open Access Hub (i.e. downloading  
data) during *last 3 months*




*Statistics of ESA Open Access Hub do not include active users downloading Sentinel data through :*

- Eumetsat (Sentinel-3)
- Partners within national collaborative ground segment (in Europe)
- Partners within international ground segment (e.g. US or Australia)

*Statistics of ESA Open Access Hub do not include active users using Sentinel data (without downloading products) through image visualisation and handling tools:*

- “EO Browser” (see next slide)



*Many Sentinel data users (in particular general public) only need basic image handling tools (i.e. not requiring data download)*

*EO Browser is a good example of such tool (~2300 unique users per week)*


<http://apps.sentinel-hub.com/eo-browser>

**Sentinel-2**  
**Forest fire in Corsica (2 January 2018)**

# ESA FDA activities – Data and Information Access Services (DIAS)

**DIAS: Creating and enabling  
a European EO Data ecosystem  
for research and business**

*A common European approach  
to EO data exploitation  
with Copernicus data at its core*




**DIAS Provider:** In charge of DIAS Back-Office Services & Interfaces operations & of providing Third-parties with the access to the DIAS computing and storage resources



- Following an Invitation to Tender issued by ESA in January 2017, **4 DIAS contracts** were signed by ESA and the respective DIAS providers on 14 December 2017 in Brussels in the presence of the EC;
- The 4 DIAS contracts were effectively kicked-off on 15 December 2017;
- The foreseen 6 months of preparatory activities (phase-in), before opening the operations, are now on going;
- The **start of operations** is planned for **June 2018**;
- **Data and service offer** form part of the DIAS concept

# DIAS ecosystem



## ALL DIAS

- offer access to *Copernicus Data and Information*, including Sentinel **data** and the Copernicus services **information**.
- may provide access to additional EO & non-EO data according to the respective DIAS business models.
- allow users (end users and Third-parties) to freely **discover, view and download available data & information**
- offer **cloud computing & storage resources** to be procured by Third-parties for further data processing
- Provide additional interfaces and services available to **store and integrate “user data” into the DIAS “data offer”**.
- Provide **interfaces for local processing**, offering standard OGC interfaces, popular cloud APIs (e.g. swift), and/or new innovative solutions (e.g. ENS, data cube, etc.) to access the Data Offer.

# ESA's contribution to ARD activities

## Current

- Sentinel-2 Surface Reflectance
- Sentinel-3
  - Land Surface Temperature
  - Synergy products
- GlobTemperature
- Proba-V

## Future

- Sentinel-1 SAR: ESA Study to develop and consolidate SAR ARD demonstrator and definition → for update see dedicated presentation in session 4
- Sentinel-2/Landsat ARD **Prototype** Products / Level-3 HS2L8
- Sentinel-2/Sentinel-3 ARD **Prototype** Products / Level-3 HS2S3

# Sentinel-2 ARD Products / Level-2A Surface Reflectance


General	Sentinel-2 Surface reflectances provided on UTM-WGS84 projection and a MRPG-based tiling grid.
External reference datasets	<ul style="list-style-type: none"><li>• DEM (PlanetDEM90) for orthorectification and topographic correction</li><li>• O<sub>3</sub> (ECMWF) for atmospheric correction</li><li>• Land Cover (CCI) for cloud masking improvement</li></ul>
Processing approach	Sen2Cor processor (Louis et al., 2017) including Dark Dense Vegetation (DDV) algorithm for aerosol optical thickness (AOT) estimation and APDA (Atmospheric Precorrected Differential Absorption) for water vapour estimation.
Input data	Level-1C products
Coverage	European coverage every 5 days since May 2017. Worldwide coverage every 5 days by mid-2018.
Spatial/temporal resolution	10/20/60m
Distribution channels	<ul style="list-style-type: none"><li>• Copernicus Open Access Hub: <a href="https://scihub.copernicus.eu/">https://scihub.copernicus.eu/</a></li></ul>



# Sentinel-2 ARD Products / Level-2A Surface Reflectance



- Systematic production over EEA39 areas started since May 2017.
- Expansion to worldwide coverage by mid 2018.



# New Sen2Cor 2.5

✓ New version of Sen2Cor released (v2.5) supporting Sentinel-2B

- Several Bugs fixed.
- Full compatibility to PSD 14.3
- Backward PSD compatibility assured.
- Improvement of Ozone selection.
- Improvement of AOT retrieval
- Improvement of Cloud Screening and Scene Classification over coastal/inland/muddy waters, urban areas/bright surfaces.



# Feedback from Sentinel-2 Validation Team on S2 L2A

# Level-2A Product



Scene Classification

AOT


WV

T33TWF  
DDV=16.0%  
Scaling=1.00  
AOT=0.214  
ECMWF=0.201  
Aeronet(AOT)=None

T33TWE  
DDV=21.1%  
Scaling=1.00  
AOT=0.178  
ECMWF=0.226  
Aeronet(AOT)=None



# Level-1C versus Level-2A Example



ESA UNCLASSIFIED - For Official Use

Susanne Mecklenburg | 06/02/2018 | Slide 30





# Level-2A Time Series



ESA UNCLASSIFIED - For Official Use

Susanne Mecklenburg | 06/02/2018 | Slide 31




# Level-2A Usage Example

## ✓ Seoul National University: Agriculture




## ✓ Sen2Coral project for coral reefs monitoring

# Level-2A Usage Example



# Level-2A performances from ACIX



MSI Band 4  
Red, 665 nm

# Sentinel-2/Landsat ARD Prototype Products

## *Level-3 HS2L8*



General	<b><u>Prototype products</u></b> HS2L8 (Harmonisation Sentinel-2 and Landsat-8) combining both missions in a single fused product on the common bands of both missions and at the spatial resolution of Sentinel-2 bands.
External reference datasets	<ul style="list-style-type: none"> <li>• DEM (PlanetDEM90) for orthorectification and topographic correction</li> <li>• O<sub>3</sub> (ECMWF) for atmospheric correction.</li> <li>• AOT(ECMWF) for atmospheric correction.</li> <li>• WV (ECMWF) for atmospheric correction.</li> </ul>
Processing approach	<ul style="list-style-type: none"> <li>• Atmospheric corrections short-term solution with SMAC (Rahman &amp; Dedieu, 1994) and mid-term solution with Sen2Cor (expanded to correct Landsat-8 data).</li> <li>• Fusion approach based on STAR-FM method (Gao et al., 2006).</li> <li>• BRDF correction (Roy et al., 2016).</li> </ul>
Input data	Sentinel-2 Level-2A and Landsat-8 Level-1T products.
Coverage	N/A
Spatial/temporal resolution	10/20/60m
Distribution channels	N/A



**Prototype products** HS2S3 (Harmonisation Sentinel-2 and Sentinel-3 OLCI) combining both missions in a single fused product on Sentinel-2 VNIR bands and using the spectral and spatial resolution of Sentinel-2 VNIR bands and the temporal resolution of Sentinel-3.

External reference datasets	<ul style="list-style-type: none"> <li>• DEM (PlanetDEM90) for orthorectification and topographic correction</li> <li>• O<sub>3</sub> (ECMWF) for atmospheric correction.</li> <li>• AOT(Aeronet) for atmospheric correction.</li> <li>• WV (ECMWF) for atmospheric correction.</li> </ul>
-----------------------------	--

Processing approach	<ul style="list-style-type: none"> <li>• Atmospheric corrections with SMAC (Rahman &amp; Dedieu, 1994).</li> <li>• Fusion approach based on STAR-FM method (Gao et al., 2006).</li> </ul>
---------------------	---

Input data	Sentinel-2 Level-1C and Sentinel-3 OLCI Level-1B.
------------	---

Coverage	N/A
----------	-----

Spatial/temporal resolution	10m
-----------------------------	-----

Distribution channels	N/A
-----------------------	-----

# Sentinel-3 Land Surface Temperature (LST)

## “quasi ARD-ready”



General	Land Surface Temperature (LST) is retrieved from thermal radiances measured by the Sea and Land Surface Temperature Radiometer (SLSTR)
External reference datasets	<ul style="list-style-type: none"> <li>• Pressure, temperature, and water vapour (ECMWF – Atmospheric HRES-10 day dataset)</li> <li>• Carbon dioxide, nitric acid, chlorofluorocarbons (CFCs) and ozone (RAMIGclim; Remedios et al, 2007)</li> <li>• Globcover Land Cover (reduced-resolution ~1 km)</li> <li>• Fractional vegetation cover climatology</li> </ul>
Processing approach	Evolution of the LST algorithm developed for the Advanced Along track Radiometer (AATSR) by CSIRO (Prata 2002) ( <i>Ref: S3-L2-SD-03-T03-ULNILU-ATBD_L2LST, 2012</i> )
Input data	11 and 12 $\mu\text{m}$ channels brightness temperature of the SLSTR nadir view
Coverage	<ul style="list-style-type: none"> <li>• Global coverage</li> </ul>
Spatial/temporal resolution	<ul style="list-style-type: none"> <li>• 1 km</li> <li>• 2 days at the equator</li> </ul>
Distribution channels	Sentinel data Hub: <a href="https://scihub.copernicus.eu/dhus/#/home">https://scihub.copernicus.eu/dhus/#/home</a>

# Sentinel-3 Synergy “quasi ARD-ready”



General	<p>Sentinel-3 Synergy products are obtained combining information from the OLCI and SRTL instruments. The standard product is a 300m SDR product.</p> <p>Moreover we provide SPOT VGT-like products in Plate Carrée projection to be consistent with standard 1 km SPOT Vegetation products consisting of instantaneous acquisition, daily and 10 day composites.</p>
External reference datasets	<ul style="list-style-type: none"> <li>• Auxiliary data: O<sub>3</sub>, Water vapour, and surface pressure (ECMWF)</li> <li>• OLCI and SLSTR Lev-1B ortho-geolocated using GETASSE30 DEM</li> </ul>
Processing approach	<ul style="list-style-type: none"> <li>• AOT estimation (North et al., 2010)</li> <li>• Level-1C resampling adopting OLCI as reference image grid</li> <li>• Currently cloud detection based on operational OLCI cloud mask</li> <li>• Ice/snow detection uses the results of the OLCI L2 Pixel classification</li> <li>• Level-2 SDR retrieval (<i>Sentinel-3 Synergy products ATBD – North, Heckel, 2010</i>)</li> </ul>
Input data	<ul style="list-style-type: none"> <li>• TOA reflectance data for the 6 solar reflective SLSTR at both nadir and forward views and 18 OLCI bands at all non-absorbing channels (bands 14,15,20)</li> </ul>
Coverage	<ul style="list-style-type: none"> <li>• Common part of the OLCI and SLSTR nadir view swaths for the daylight part of the SENTINEL-3 orbit, i.e. the entire OLCI swath: ~1 270 km. Two days at the equator</li> </ul>
Spatial/temporal resolution	<ul style="list-style-type: none"> <li>• 300 m for Synergy L2 SDR and 1 km for VGT-like products (Daily and 10 day)</li> </ul>
Distribution channels	<p>Sentinel data Hub: <a href="https://scihub.copernicus.eu/dhus/#/home">https://scihub.copernicus.eu/dhus/#/home</a> (Currently only Exp. Hub)</p>

# Globe Temperature “quasi ARD-ready”



General	The aim of ESA's GlobTemperature project is to promote the wider uptake of global-scale satellite LST, to merge LST data set from thermal infrared (geostationary & polar orbiter) and passive microwave satellite data to provide best possible coverage, including information on the diurnal cycle and clear-sky bias.
External reference datasets	All datasets used at single sensor level for generating L2-LST products
Processing approach	<ul style="list-style-type: none"> <li>• Level-2: geophysical variables derived from the Level 1b source data from a single orbit of a sensor at the same resolution and location as the Level 1b data</li> <li>• Level-3: “uncollated” – single sensor data regridded and/or spatially averaged onto a spatial grid without combination of observations from overlapping orbits; “collated” - multiple orbits from a single sensor have been combined onto a spatial/temporal grid</li> <li>• Level-4: datasets constructed from multiple orbits and sensor</li> <li>• Complete description of all processing variables at L2-4 in <i>GlobT-WP3-DEL-11, 2018</i> (Globtemp Product user Guide)</li> </ul>
Input data	Land Surface Temperature (LST) data sets from: (A)ATSR; SEVIRI; SSM/I; AMSR-E, MODIS, JAMI, GOES, MTSAT, ATSR-CDR; SLSTR
Coverage	<ul style="list-style-type: none"> <li>• Global coverage</li> </ul>
Spatial/temporal resolution	<ul style="list-style-type: none"> <li>• 1 km for merged dataset</li> <li>• 3 hourly daily for merged dataset</li> </ul>
Distribution channels	GlobTemperature Project portal: <a href="http://data.globtemperature.info/">http://data.globtemperature.info/</a>

ES

Susanne Mecklenburg | 06/02/2018 | Slide 41



European Space Agency

General	Proba-V Top-Of-Canopy (TOC) Reflectances provided as synthesis products in Plate Carrée projection for different compositing period: 1-day, 5-days (for 100m), 10-days [ <i>Specs in continuity with SPOT-VGT data record</i> ]
External reference datasets	<ul style="list-style-type: none"> <li>• DEM (GLSDEM) for horectification and surf pressure</li> <li>• WV (ECMWF) and O3 (TOMS climatology) for A/C</li> <li>• Surface Reflectance climatology (MERIS) and Land Cover (CCI) for cloud masking</li> </ul>
Processing approach	<ul style="list-style-type: none"> <li>• Cloud/Shadow/Snow masking (<i>Wolters et al. 2017</i>)</li> <li>• AOT estimation (<i>Maisongrande et al., 2004</i>)</li> <li>• A/C using SMAC (<i>Rahman et al., 1994</i>)</li> <li>• Compositing using MVC (<i>Holben, 1986</i>)</li> </ul>
Input data	<ul style="list-style-type: none"> <li>• L2-B (intermediate product, not for users): atmospherically corrected segment files to be combined in the synthesis products</li> </ul>
Coverage	<ul style="list-style-type: none"> <li>• Global quasi-daily coverage, two days for [-35°:35°]</li> </ul>
Spatial/temporal resolution	<ul style="list-style-type: none"> <li>• 1km and 333m with quasi-daily revisit</li> <li>• 100 m 5-days revisit</li> </ul>
Distribution channels	<ul style="list-style-type: none"> <li>• VITO PDF portal: <a href="http://www.vito-eodata.be/PDF/">http://www.vito-eodata.be/PDF/</a></li> <li>• Proba-V MEP: <a href="https://proba-v-mep.esa.int">https://proba-v-mep.esa.int</a></li> </ul>



# ESA's contribution to ARD - outlook




- **“Quasi” compliant to ARD** Product Family Specifications
- **DOI planned** for product type (e.g. LST) and in future also algorithm (e.g. LST retrieval) identification
- **New ARD activities**
  - Prototype products over land being developed
  - ESA Study to develop and consolidate SAR ARD demonstrator and definition
- **On-going evaluation for the Evolution of the Digital Elevation Model (DEM)** used in the Copernicus Space Component Data Processing; ESA performing study to provide recommendations for the evolution of the DEM to be used for the generation of Copernicus Space Component (CSC) core products:
  - Collect DEM requirements from the different CSC elements (i.e. Sentinel missions and Copernicus Contributing Missions).
  - Identify different candidate DEM available.
  - Provide preliminary quantitative assessment of the impact of the pre-selected DEMs on the different missions' products data quality and on the production operational set-up.
  - Financial assessment (cost, license, maintenance ...).
  - Final conclusions and recommendations for the choice of the DEMs (by April 2018).



## Day 3: Thursday, 1 March 2018

### Session 3: Harmonization and Consistency

09:00 – 09:30	<b>Keynote: Reaching new users through the CEOS Analysis Ready Data for Land (CARD4L) Framework</b>	<b>Steven Hosford, ESA, Italy</b>
09:30 – 09:50	Quality assessment of the Level-2 Sentinel-2 products through the Harmonized Landsat and Sentinel-2 (HLS) data set	Martin Claverie, UCLouvain, Belgium
09:50 – 10:10	Towards achieving continental scale field validation and multi-sensor interoperability of satellite derived surface reflectance in Australia	Medhavy Thankappan, Geoscience Australia, Australia
10:10 – 10:30	Validating PCI ARD data with Sen2Cor and LaSRC including BRDF correction and Topographic Normalization.	Wolfgang Lück, Pci Geomatics, Canada
10:30 – 10:50	A new 'system state vector' of optical, thermal and microwave EO data for terrestrial change detection	Mathias Disney, UCL Geography, UK



# Take home messages



- The **Copernicus Space Component delivers high quality and timely data** to the user community
  - Current fleet of missions: nominal operations and all core data products released
  - C/D units in preparation
  - Long-term Scenario/ future Sentinel extension and expansion/ consolidated
- **ESA's provides important contribution to FDA activities**
  - Excellent Copernicus data access service provided, including data provision working in partnership in international environment
  - DIAS contracts now up and running
- **ESA's data products over land** provide a considerable contribution to the CEOS ARD portfolio and are **"quasi" compliant to ARD** principles
- **New ARD activities triggered by ESA**
- **ESA lead for MRI activities has been appointed:** Ferran Gascon (Sentinel-2 mission manager)