



CENTRE NATIONAL D'ÉTUDES SPATIALES

CNES in-orbit calibration activities for visible and NIR sensors

Patrice HENRY
CNES – Toulouse

CNES background in calibration activity

- **CNES has developed different calibration methods over natural targets for visible and NIR optical sensors**
 - ◆ Rayleigh scattering over ocean
 - ◆ Sun glint over ocean
 - ◆ Deep convective clouds (DCC)
 - ◆ Stable African deserts
 - ◆ Antarctica (Dome C site)
 - ◆ Autonomous calibration station (for high resolution)
 - ◆ Lunar calibration
- **Most of them are used on an operational basis**
 - ◆ Monitoring the CNES sensors calibration (SPOT, VGT, POLDER...)
 - ◆ Inter calibration with other sensors (MERIS, SeaWiFS, AVHRR, MODIS, Formosat2, Kompsat2...)

Calibration monitoring activities for CNES missions (1)

- **POLDER 3 on-board PARASOL (A-train)**
 - ♦ Routine calibration monitoring using DCC calibration (every month)
 - ♦ Regular calibration check with other methods : Rayleigh scattering, sun glint, desert sites (every 6 months)
- **VGT2 on-board SPOT5**
 - ♦ Misfunctioning of the on-board calibration device
 - ♦ Routine calibration monitoring using desert sites (every month)
 - ♦ Regular calibration check with other methods : Rayleigh scattering, sun glint, Antarctica sites (once a year)
- **High resolution instruments on-board SPOT4 & SPOT5**
 - ♦ Routine calibration monitoring over desert sites
 - ♦ Calibration campaigns over La Crau and Negev desert (twice a year)

Calibration monitoring activities for CNES missions (2)

- **IIR on-board CALIPSO (A-Train)**
 - ♦ Calibration monitoring using on-board device
- **IASI on-board METOP**
 - ♦ Radiometric and spectral calibration
 - ♦ Regular performance checks
 - ♦ IASI / AIRS inter-calibration

Soon to come :

- **Recalibration of VGT1 data**
 - ♦ To insure with VGT2 a consistent set of data over 10 years
- **Preparation of Pleiades and Venus in-orbit calibration**

Cal/Val activities in cooperation with other agencies

■ FORMOSAT2 calibration for NSPO

- ♦ Activity started in 2005
- ♦ Calibration monitoring over deserts, absolute calibration over La Crau, programming gains optimization using the worldwide CNES data base

■ Kompsat2 Cal/Val activities for KARI

- ♦ Activity in progress
- ♦ Geometric calibration, FTM assessment, radiometric calibration...

■ MERIS for ESA

- ♦ Desert, Rayleigh and glitter calibration on an operational basis


■ Other inter-calibration activities

- ♦ MODIS, AVHRR (in the framework of GSICS activities)

Recent results


2 years of PARASOL calibration over DCC (1)

- Calibration measurements performed every month (using 1 week of data)




2 years of PARASOL calibration over DCC (2)

- Calibration model (fitted with 18 months of measurements) + extrapolation performed every month using new measurements



VGT2 calibration monitoring over desert sites


VGT2/B0 calibration




MERIS calibration monitoring

based on cross-calibration with PARASOL over 20 desert sites


490




665



560




865



No significant variation with time



Validation of MERIS calibration using Rayleigh scattering and sun glint methods



No discrepancy greater than 2%

Formosat 2 calibration monitoring

based on cross-calibration with POLDER1 over 20 desert sites



Spectral band	Loss (in 2.5 years)
B1	~ 4 %
B2	~ 1.5 %
B3	< 1 %
B4	negligeable
PAN	< 1 %

